

Mini Review

Open Access

Arachidonic acid in Alzheimer's disease

Mélanie H. Thomas¹, Sandra Pelleieux^{1,2}, Nicolas Vitale³ and Jean Luc Olivier^{1,2*}¹Unité de Recherche Aliment et Fonctionnalité des Produits Animaux (URAFPA), INRA USC 0340, Université de Lorraine, Nancy, France²Service de Biochimie-Biologie Moléculaire, Hôpital Central, CHU de Nancy, Nancy, France³Institut des Neurosciences Cellulaires et Intégratives (INCI), UPR CNRS 3212, Université de Strasbourg, Strasbourg, France

Article Info

Article Notes

Received: October 8, 2016

Accepted: December 12, 2016

*Correspondence:

Dr. Jean Luc Olivier, Laboratoire de Biochimie – Biologie Moléculaire, Hôpital Central, 29 avenue du Maréchal De Lattre de Tassigny, 54035 NANCY Cedex, (Tel : 33 (0) 3 83 85 27 85 (secrétariat) / Fax /33 (0)3 83 85 19 69, Email: jl.olivier@chu-nancy.fr

© 2016 Olivier JL. This article is distributed under the terms of the Creative Commons Attribution 4.0 International License

ABSTRACT

Alzheimer's disease is a very complex disease in which neuroinflammation and synaptic dysfunctions play a critical role in association with the two well-known molecular agents of the disease, the $A\beta_{1-42}$ peptide oligomers and the hyperphosphorylated tau protein. Arachidonic acid, the main member of the ω -6 series, is quantitatively the second polyunsaturated fatty acid in brain and is mainly esterified in membrane phospholipids. It is specifically released by the cytosolic phospholipase A_2 whose inhibition or gene suppression counteract the deleterious effects of $A\beta_{1-42}$ peptide oligomers on cognitive abilities. Arachidonic acid can be reincorporated under the action of the acyl-CoA synthetase 4 and lysophospholipid acyltransferases which remain to be characterized. Free arachidonic acid can be involved in Alzheimer's disease through several mechanisms. First it is converted by cyclooxygenases-1/2 and the specific prostaglandin synthases into PGE2 and PGD2 which contributes to the occurrence and progression of neuroinflammation. Neuroinflammation has positive as well as negative effects, by favoring $A\beta_{1-42}$ peptide clearance on one hand and by increasing the production of neurotoxic compounds on the other hand. Second, free arachidonic acid is also involved in synaptic functions as a retrograde messenger and as a regulator of neuromediator exocytosis. Third, some studies indicated that free arachidonic acid and its derivatives activate kinases involved in tau hyperphosphorylation. In addition, the dietary intakes of arachidonic acid in western food increased in the last period. Taken together, these various reports support the hypothesis that arachidonic acid is interesting target in nutrition-based preventive strategies against this disease.

Introduction

All the recent clinical trials against Alzheimer's disease (AD) failed to evidence any efficiency against the sporadic or late onset cases. A better understanding of AD mechanisms is required to open new therapeutic or preventive strategies. Lipids are important actors in AD as shown by the numerous works devoted to the putative neuroprotective role of docosahexaenoic acid (DHA, C22:6 ω -3), the main polyunsaturated fatty acid (PUFA) in brain¹. By contrast, the role of arachidonic acid (ARA, C20:4 ω -6) was less extensively studied, despite the fact that it is the second PUFA in the brain. ARA corresponds to around 20% of the total amount of the neuronal fatty acids and is mainly esterified in membrane phospholipids. After its release by phospholipases A_2 , ARA is converted by several enzymes into numerous eicosanoids which are actors of neuroinflammation². But ARA can also be directly involved in synaptic functions as a free fatty acid³. Therefore, the level of intracellular free ARA and the balance between the releasing enzymes and those which allows its incorporation in membrane phospholipids can be critical for AD-

associated phenomena such as neuroinflammation and synaptic dysfunctions. Both these events can be observed in AD murine models before the two historical pathognomonic AD signs, the amyloid plaques and the neurofibrillary tangles which are respectively formed by the two known AD agents, Aβ peptide and hyperphosphorylated tau. Finally, western food which contains excessive ω-6 / ω-3 ratios⁴ (above the recommended value of 5) might favor its incorporation in brain lipids and its influence on AD mechanisms. As a consequence, a better knowledge of the relationship between ARA and these mechanisms is useful for the therapy or prevention of AD. These different points will be discussed in this mini-review.

Mobilization of arachidonic acid in neuronal and glial cells

Free ARA is released from membrane phospholipids by

several phospholipases A₂ which constitute a superfamily including intracellular as well as secretory and Ca²⁺-dependent or -independent enzymes⁵. Group IVA PLA₂ (or cPLA₂-α) is the only PLA₂ which specifically release ARA^{6,7}. This cytosolic enzyme is activated and translocated to membranes after a rise of cytosolic calcium concentration and the phosphorylation of its Ser⁵⁰⁵ by the MAP-kinase cascade⁷ (Figure 1, left). The involvement of cPLA₂-α in AD was first evidenced by its higher expression in AD patient brain⁸. Several works also showed that cPLA-α mediates the effects of Aβ₁₋₄₂ peptide in astrocytes and microglial cells^{9,10}. cPLA₂-α is also expressed in neuronal cells^{11,12} which are protected from the pro-apoptotic effects of Aβ₁₋₄₂ peptide oligomers by the reduction of its activity or expression¹¹. Aβ₁₋₄₂ peptide oligomers activate cPLA₂-α through the stimulation of MAP kinase cascade (Figure 1, left) since Erk1/2 inhibitors suppress the




Figure 1: Pleiotropic effects of arachidonic acid in Alzheimer’s disease:

Left: arachidonic acid (ARA) is released by cytosolic phospholipase A₂ (cPLA₂-α) in neuronal cells and glial cells. It is re-incorporated in phospholipids after its conversion into arachidonyl-CoA by acyl-CoA synthetase 4 (ACSL4). Free ARA influences synaptic function by acting as retrograde messenger and by opening Kv channels which lead to exocytosis of neuromediator and induction of LTP. **Top:** arachidonic acid activates several kinases which directly or indirectly contribute to increase tau phosphorylation levels. **Right:** free Arachidonic acid is converted into PGG₂ by cyclooxygenase-2 (COX-2) which is overexpressed in neuronal cells in Alzheimer’s disease. PGG₂ is converted in PGD₂ and PGE₂ by corresponding prostaglandin synthases in astrocytes and microglial cells. PGE₂ bind to EP₂ and EP₄ receptors on microglial cell membranes. These receptors have different effects in neuroinflammation, stimulation of Aβ clearance or production of neurotoxic molecules. These positive or negative effects illustrate the dual role of neuroinflammation.

cPLA₂- α phosphorylation. Several signaling pathways downstream of cPLA₂- α activation have been characterized in neuronal cells such as stimulation of sphingomyelinase¹³ or PLD2¹⁴ activities. Finally, the critical role of cPLA₂- α was definitively established by *in vivo* experiments showing that its gene suppression leads to resistance to A β peptide in a transgenic murine line bearing a double mutation of human APP¹² or in wild-type mice submitted to intracerebroventricular injections of A β peptide¹⁵. But the way by which cPLA₂- α -released ARA mediates A β ₁₋₄₂ oligomer neurotoxicity remains to be identified.

Free ARA is incorporated into membrane phospholipids through two steps, 1) its esterification with Coenzyme A into arachidonyl-CoA by acyl-CoA synthetases (ACSLs), 2) its transfer from arachidonyl-CoA to lysophospholipids by various acyl-CoA lysophospholipids acyltransferases (LPATs). Among the various ACSLs, ACSL4 specifically esterifies ARA (Figure 1, left) and was first described in steroidogenic tissues¹⁶. In addition to the ubiquitous isoform, a neuronal specific isoform was identified and differs from the first one by an additional 41 amino-acid N-terminal domain¹⁷. Mutant ACSL4 gene was associated to X-linked mental retardation¹⁸ and has been involved in neuronal differentiation, axonal transportations and synapse formation^{17,18,19}. Channeling of ARA by ACSL4 and Lysophosphatidylinositol-acyltransferase-1 (LPIAT1) to phosphatidylinositol (PI) has been described in CHO cells²⁰. But the role of the various LPATs in the ARA incorporation in brain phospholipids remains to be characterized in more details. ACSL4 and LPATs could reduce the free ARA level in neuronal and glial cells, thus limiting the impact of enhanced cPLA₂- α activity on this level and the consequences on neuroinflammation and synapse dysfunctions.

Neuroinflammation and AD: role of arachidonic acid and its derivatives

Neuroinflammation plays a positive role in the early AD steps by contributing to A β peptide clearance but can contribute to synapse alterations in the advanced stage of the disease when A β peptide accumulates and aggregates in amyloid plaques²¹. Eicosanoids formed from free ARA are key mediators in neuroinflammation. Among the various enzymes which produced these eicosanoids, cyclooxygenases-1 and -2 (COX-1/COX-2) convert ARA into PGH₂²². Then, PGH₂ is used by various prostaglandin synthases to produce the various prostaglandins in the different tissues²³. Nonsteroidal anti-inflammatory drugs (NSAIDs) are well known inhibitors of COX-1 and COX-2. Their involvement in AD has first been suggested by studies which evidenced a reduction of AD risk in patients treated by NSAIDs²⁴⁻²⁶. In the majority of organs and tissues, COX-1 is constitutively expressed and COX-2 gene expression is induced especially in case of inflammation. On the contrary, brain tissues and especially neuronal cells constitutively

express COX-2 in basic condition²⁷ (Figure 1, left). But COX-2 overexpression in brain of AD patients has been reported and correlated to the progression of the disease in several studies^{28,29}. Since A β ₁₋₄₂ oligomers do not induce the COX-2 expression in human microglial cell primary cultures³⁰, the COX-2 overexpression observed in AD patient brains is likely to be generated in neuronal cells. On the contrary to COX-2, COX-1 is overexpressed by microglial cells around amyloid plaques³¹. In addition, the expression of COX-3, a splice variant of COX-1, has also been reported in AD hippocampus and A β ₁₋₄₂ treated neuronal primary culture neuronal cells³². PGH₂ can be secreted by neuronal cells and converted into other prostaglandins by glial cells. In the brain, the most abundant prostaglandins are PGD₂ and PGE₂ (Figure 1, left), which are synthesized by several PGD₂ and PGE₂ synthases. Expression levels of hematopoietic PGD₂ synthase (one of the two PGD₂ synthases expressed in the brain) are enhanced in microglial cells and astrocytes of AD transgenic mice and patients³³. As a result PGD₂ is overproduced in these glial cells surrounding the amyloid plaques³³. PGE₂ binds to several receptors and have opposite effects on inflammation depending on the receptor type. EP₂ receptor mediates negative inflammatory effects (Figure 1, right) since its gene suppression reduces the microglial toxic inflammatory response and restored A β clearance³⁴. By contrast, suppression of the EP₄ receptor increases the inflammatory response in the brain of the APPSwe-PS1 Δ E9 mouse³⁵.

Five-lipoxygenase (5-LOX) converts free ARA into 5-hydroxyperoxyeicosatetraenoic acid [5-HPETE], which is then stabilized into 5-hydroxyeicosatetraenoic acid (5-HETE) or converted by 5-LOX into leukotriene A₄ (LTA₄). 5-LOX gene deletion attenuates the worsening effect of LPS-induced neuroinflammation in transgenic AD-model mice³⁶. But the use of 5-LOX inhibitors demonstrated that 5-LOX and its products have a stronger impact on A β peptide production and tau phosphorylation in neuronal cells³⁷ than in the proliferation of glial cells and neuroinflammation.

Despite the data of retrospective epidemiologic studies which reported a lower AD risk in NSAIDs-treated populations, randomized controlled trial failed to support the efficiency of these drugs in the AD therapy^{26,38}. Therefore, other approaches are needed to explain the discrepancy between the data of the observational studies or the works on AD animal models on one hand and those of the clinical trials on the other hand. Parallel mechanisms could explain the role of COX inhibitors in AD. For example, higher plasma concentrations of tryptophan metabolites were found in AD patients suggesting the involvement of the kynurenine pathway which influences the serotonergic and glutamatergic neurotransmission³⁹. Ibuprofen drastically reduced the expression of the neuronal

tryptophan 2,3-dioxygenase (Tdo2), which encodes an enzyme that metabolizes tryptophan to kynurenine while Tdo2 inhibition prevented behavioural deficits in the AD model APPSwe-PS1ΔE9 mice⁴⁰. However, the influence of ARA and its metabolites on Tdo2 expression remains to be established.

Arachidonic acid and synaptic functions

Before its deposit in amyloid plaques, Aβ₁₋₄₂ peptide forms oligomers which are now recognized as the main pathological actor in the early steps of AD. These oligomers initially alter the remodeling of the synaptic network and memory abilities by inhibiting long-term potentiation (LTP)⁴¹. Free ARA produced by the postsynaptic neuron acts as a retrograde messenger which increases the neurotransmitter release by the presynaptic neuron and results in the induction of LTP⁴² (Figure 1, left). Regulation of the pre and post-synaptic Kv channels by ARA is also involved in its effect on LTP⁴³. Moreover the LTP was reduced by cPLA₂-α gene suppression or its inhibition by arachidonyl trifluoromethyl ketone (AACOCF3)⁴⁴. Thus ARA seems to be necessary for the LTP induction.

Free ARA activates the soluble N ethylmaleimide-attachment receptors (SNARE), which are required for the fusion of synaptic vesicles with the plasma membrane⁴⁵ (Figure 1, left). Vesicle SNARE protein VAMP-2 interacts with the plasma membrane SNARE proteins SNAP25 and syntaxin-1 to form a complex which promotes membrane fusion. ARA induces the binding of syntaxin-1 to the SNARE complex in presence of Munc18-1, which is a critical regulator of this process⁴⁶. The involvement of ARA in the vesicle membrane fusion was also demonstrated by the formation of TIP30 complex which associates Tip30 endophilin B and ACSL4⁴⁷. In this complex, TIP30 transfers ACSL4-produced arachidonyl-CoA onto phosphatidic acid and thus forms new species which induce close contact between membranes. Initial work on neuroendocrine cells revealed increased ARA production during exocytosis and clearly showed that ARA provision facilitated vesicular release⁴⁸. Taken together, these studies support the hypothesis that modulation of free ARA levels can impact synaptic functions. Excessive maintenance of LTP or release of neurotransmitter could result from excessive levels of free ARA levels and have detrimental effects on memory storage.

Arachidonic acid and tau

Tau is a microtubule associated protein which is physiologically weakly phosphorylated and concentrated in the neuronal axons⁴⁹ (Figure, top). The C-terminal domain of Tau binds to α and β-tubulin to assemble microtubules and regulates the axonal transport of neurotransmitters. In AD, an excessive kinase activity and/or a shortfall of phosphatase activity lead to Tau hyperphosphorylation⁴⁹. Hyperphosphorylated Tau accumulates in the dendrites

and cell body of the neuron, forms helical filaments and subsequently neurofibrillary tangles⁵⁰.

Several studies reported that free ARA *in vitro* induced tau polymerization. The ARA concentration required for the polymerization differs according to the various tau isoforms which are generated by alternative splicing⁵¹. These polymers activate microglial cells⁵².

Among the various kinases which phosphorylate tau, two enzymes, PKNα and PKCξ bind and are activated by ARA^{53,54}. PKNα is one of the 3 PKC-homologous serine/protein kinases (PKN), is highly expressed in brain⁵⁴ and accumulates in neurofibrillary tangles^{55,56}. Increased expression levels of PKCξ, an atypical Protein Kinase C family member, have been measured in neuronal membrane fraction of AD TG2576 mice⁵⁷ and in T-lymphocytes of patients affected by severe forms of AD⁵⁸. PKCξ contributes to tau phosphorylation by targeting leucine-rich repeat kinase 2 [LRRK2] whose mutation is the most frequent cause of the genetic forms of Parkinson disease⁵⁹.

Lipoxygenase-generated derivatives of ARA are also involved in tau phosphorylation. Suppression of 5-LOX gene expression improves Tau and Aβ-related mechanisms in AD mouse models⁶⁰. In addition, the dual enzyme 12/15-lipoxygenase and its products, 12(S)-HETE and 15(S)-HETE modulates Tau metabolism specifically via the cdk5 kinase pathway⁶¹.

Conclusion

Free ARA contributes to AD progression through various mechanisms. Through its conversion into pro-inflammatory eicosanoids, it participates to neuroinflammation. ARA is directly involved in synaptic functions as a retrograde messenger and a regulator of neuromediator exocytosis. Finally, some works also indicate that ARA might have some influence on tau phosphorylation and polymerization. All these data shows that ARA has pleiotropic effects in AD and might be an interesting target in the fight against this complex disease.

ARA is provided by diet, directly from animal products and indirectly from the conversion of linoleic acid found in the vegetal oils. Linoleic acid and ARA amounts increased in the last 40 years period in the western diets. Therefore the influence of dietary ARA on the occurrence of AD is an important issue for the prevention of the disease. Unfortunately, only two studies were performed on transgenic AD murine model to measure the impact of dietary ARA on the pathological process and found opposite results on Aβ production and deposition^{62,63}. Therefore, additional studies are necessary to clarify the dietary ARA impact in AD and to identify the underlying mechanisms, keeping in mind the perspective of nutrition-based preventive strategies of AD.

Acknowledgements

We thank the French France-Alzheimer association and the French region of Lorraine for the financial support to our work on the role of dietary arachidonic acid in Alzheimer's disease which was the initiation point of this minireview.

References

- Luchtman D. W. and Song C. Cognitive enhancement by omega-3 fatty acids from child-hood to old age findings from animal and clinical studies. *Neuropharmacology*. 2013; 64: 550-565.
- Funk C. D. Prostaglandins and leukotrienes advances in eicosanoid biology. *Science*. 2001; 294(5548): 1871-1875.
- Latham C F, Osborne S L, Cryle M J et al. Arachidonic acid potentiates exocytosis and allows neuronal SNARE complex to interact with Munc18a. *J Neurochem*. 2007; 100(6): 1543-1554.
- Rett B S, Whelan J. Increasing dietary linoleic acid does not increase tissue arachidonic acid content in adults consuming Western type diets a systematic review. *Nutr Metab*. 2011; 8: 36.
- Burke JE, Dennis EA. Phospholipase A2 structure/function mechanism and signaling. *J Lipid Res*. 2009; 50 Suppl: S237-42.
- Clark JD, Lin LL, Kriz RW, et al. A novel arachidonic acid-selective cytosolic PLA2 contains a Ca(2+)-dependent translocation domain with homology to PKC and GAP. *Cell*. 1991; 65(6): 1043-51.
- Leslie CC. Cytosolic phospholipase A₂: physiological function and role in disease. *J Lipid Res*. 2015; 56(8): 1386-402.
- Stephenson DT, Lemere CA, Selkoe, DJ, et al. Cytosolic phospholipase A2 (cPLA2) immunoreactivity is elevated in Alzheimer's disease brain. *Neurobiol Dis*. 1996; 3(1): 51-63.
- Hicks JB, Lai Y, Sheng W, et al. Amyloid-beta peptide induces temporal membrane biphasic changes in astrocytes through cytosolic phospholipase A2. *Biochim Biophys Acta*. 2008; 1778(11): 2512-9.
- Szaingurten-Solodkin I, Hadad N, Levy R. Regulatory role of cytosolic phospholipase A2alpha in NADPH oxidase activity and in inducible nitric oxide synthase induction by aggregated Abeta1-42 in microglia. *Glia*. 2009; 57(16): 1727-40.
- Kriem B, Sponne I, Fifre A, et al. Cytosolic phospholipase A2 mediates neuronal apoptosis induced by soluble oligomers of the amyloid-beta peptide. *FASEB J*. 2005; 19(1): 85-7.
- Sanchez-Mejia RO, Newman JW, Toh S, et al. Phospholipase A2 reduction ameliorates cognitive deficits in a mouse model of Alzheimer's disease. *Nat Neurosci*. 2008; 11(11): 1311-8.
- Malaplate-Armand C, Florent-Bécharde S, Youssef I, et al. Soluble oligomers of amyloid-beta peptide induce neuronal apoptosis by activating a cPLA2-dependent sphingomyelinase-ceramide pathway. *Neurobiol Dis*. 2006; 23(1): 178-89.
- Oliveira TG, Chan RB, Tian H, et al. Phospholipase d2 ablation ameliorates Alzheimer's disease-linked synaptic dysfunction and cognitive deficits. *J Neurosci Off J Soc Neurosci*. 2010; 30(49): 16419-28.
- Desbène C, Malaplate-Armand C, Youssef I, et al. Critical role of cPLA2 in Aβ oligomer-induced neurodegeneration and memory deficit. *Neurobiol Aging*. 2012; 33: 1123 e17-29.
- Kang MJ, Fujino T, Sasano H, et al. A novel arachidonate-preferring acyl-CoA synthetase is present in steroidogenic cells of the rat adrenal; ovary; and testis. *Proc Natl Acad Sci U S A*. 1997; 94(7): 2880-4.
- Cho, YY. A novel role of brain-type ACS4 isotype in neuronal differentiation. *Biochem Biophys Res Commun*. 2012 ; 419(3): 505-10.
- Meloni I, Parri V, De Filippis R, et al. The XLMR gene ACSL4 plays a role in dendritic spine architecture. *Neuroscience*. 2009; 159(2): 657-69.
- Liu Z, Huang Y, Zhang Y, et al. Drosophila Acyl-CoA synthetase long-chain family member 4 regulates axonal transport of synaptic vesicles and is required for synaptic development and transmission. *J Neurosci*. 2011; 31(6): 2052-63.
- Küch EM, Vellaramkalayil R, Zhang I, et al. Differentially localized acyl-CoA synthetase 4 isoenzymes mediate the metabolic channeling of fatty acids towards phosphatidylinositol. *Biochim Biophys Acta*. 2013; 1841(2): 227-239.
- Heneka MT, Carson MJ, El Khoury J, Landreth GE, Brosseron F, Feinstein DL, et al. Neuroinflammation in Alzheimer's disease. *Lancet Neurol*. 2015; 14: 388-405
- Smith WL, Song I. The enzymology of prostaglandin endoperoxide H synthases 1 and 2. *Prostaglandins Other Lipid Mediat*. 2002; 68 69: 115 28.
- Pecchi E, Dallaporta M, Jean A, et al. Prostaglandins and sickness behavior old story new insights. *Physiol Behav*. 2009; 97(3 4): 279 92.
- Szekely CA, Zandi PP. Non-steroidal anti-inflammatory drugs and Alzheimer's disease the epidemiological evidence. *CNS Neurol Disord Drug Targets*. 2010; 9(2): 132-9.
- Côté S, Carmichael PH, Verreault R, et al. Nonsteroidal anti-inflammatory drug use and the risk of cognitive impairment and Alzheimer's disease. *Alzheimers Dement*. 2012 May; 8(3): 219-26.
- Wang J, Tan L, Wang HF, et al. Anti-inflammatory drugs and risk of Alzheimer's disease, an updated systematic review and meta-analysis. *J Alzheimers Dis*. 2015; 44(2): 385-96.
- Minghetti, L. Cyclooxygenase-2 [COX-2] in inflammatory and degenerative brain diseases. *J Neuropathol Exp Neurol*. 2004; 63(9): 901-10.
- Ho L, Purohit D, Haroutunian V, et al. Neuronal cyclooxygenase 2 expression in the hippocampal formation as a function of the clinical progression of Alzheimer disease. *Arch Neurol*. 2001; 58(3): 487 92.
- Fujimi K, Noda K, Sasaki K, et al. Altered expression of COX-2 in subdivisions of the hippocampus during aging and in Alzheimer's disease the Hisayama Study. *Dement Geriatr Cogn Disord*. 2007; 23(6): 423 31.
- Hoozemans JJ, Veerhuis R, Janssen I, et al. The role of cyclo-oxygenase 1 and 2 activity in prostaglandin E secretion by cultured human adult microglia, implications for Alzheimer's disease. *Brain Res*. 2002; 951(2): 218-26.
- Yermakova AV, Rollins J, Callahan LM, et al. Cyclooxygenase-1 in human Alzheimer and control brain quantitative analysis of expression by microglia and CA3 hippocampal neurons. *J Neuropathol Exp Neurol*. 1999; 58(11): 1135-46.
- Cui JG, Kuroda H, Chandrasekharan NV, et al. Cyclooxygenase-3 gene expression in Alzheimer hippocampus and in stressed human neural cells. *Neurochem Res*. 2004; 29(9): 1731-7.
- Mohri I, Kadoyama K, Kanekiyo T, et al. Hematopoietic prostaglandin D synthase and DP1 receptor are selectively upregulated in microglia and astrocytes within senile plaques from human patients and in a mouse model of Alzheimer disease. *J Neuropathol Exp Neurol*. 2007; 66(6): 469 80.
- Johansson JU, Woodling NS, Wang Q, et al. Prostaglandin signaling suppresses beneficial microglial function in Alzheimer's disease models. *J Clin Invest*. 2015; 125(1): 350 64.
- Woodling NS, Wang Q, Priyam PG, et al. Suppression of Alzheimer-associated inflammation by microglial prostaglandin-E2 EP4 receptor signaling. *J Neurosci Off J Soc Neurosci*. 2014; 34(17): 5882 94.
- Joshi YB, Giannopoulos PF, Chu J, et al. Modulation of lipopolysaccharide-induced memory insult γ-secretase and neuroinflammation in triple transgenic mice by 5-lipoxygenase. *Neurobiol Aging*. 2014; 35(5): 1024-31.

37. Chu J, Li JG, Praticò D. Zileuton improves memory deficits amyloid and tau pathology in a mouse model of Alzheimer's disease with plaques and tangles. *PLoS One*. 2013; 8: e70991.
38. Miguel-Álvarez M1, Santos-Lozano A, Sanchis-Gomar F, et al. Non-steroidal anti-inflammatory drugs as a treatment for Alzheimer's disease a systematic review and meta-analysis of treatment effect. *Drugs Aging*. 2015; 32(2): 139-47.
39. Gulaj E, Pawlak K, Bien B, et al. Kynurenine and its metabolites in Alzheimer's disease patients. *Adv Med Sci*. 2010; 55(2): 204-11.
40. Woodling NS, Colas D, Wang Q, et al. Cyclooxygenase inhibition targets neurons to prevent early behavioural decline in Alzheimer's disease model mice. *Brain*. 2016; 139(Pt 7): 2063-81.
41. Shankar GM, Li S, Mehta TH, et al. Amyloid-beta protein dimers isolated directly from Alzheimer's brains impair synaptic plasticity and memory. *Nat Med*. 2008; 14(8): 837-42.
42. Nishizaki T, Nomura T, Matsuoka T, et al. Arachidonic acid as a messenger for the expression of long-term potentiation. *Biochem Biophys Res Commun*. 1999; 254(2): 446-9.
43. Angelova PR, Müller WS. Arachidonic acid potently inhibits both postsynaptic-type Kv4.2 and presynaptic-type Kv1.4 IA potassium channels. *Eur J Neurosci*. 2009; 29(10): 1943-50.
44. Su LD, Wang DJ, Yang D, et al. Retrograde cPLA2 α /arachidonic acid/2-AG signaling is essential for cerebellar depolarization-induced suppression of excitation and long-term potentiation. *Cerebellum*. 2013; 12(3): 297-9.
45. Jahn R, Scheller RH. SNAREs--engines for membrane fusion. *Nat Rev Mol Cell Biol*. 2006; 7(9): 631-43.
46. Latham CF, Osborne SL, Cryle MJ, et al. Arachidonic acid potentiates exocytosis and allows neuronal SNARE complex to interact with Munc18a. *J Neurochem*. 2007; 100(6): 1543-54.
47. Zhang C, Li A, Gao S, et al. The TIP30 protein complex, arachidonic acid and coenzyme A are required for vesicle membrane fusion. *PLoS One*. 2011; 6(6): e21233.
48. Vitale N, Thiersé D, Bader MF. Melittin promotes exocytosis in neuroendocrine cells through the activation of phospholipase A₂. *Regul Pept*. 2010; 165(1): 111-6.
49. Khan SS, Bloom GS. Tau The Center of a Signaling Nexus in Alzheimer's Disease. *Front Neurosci*. 2016; 9: 10-31.
50. Billingsley ML, Kincaid RL. Regulated phosphorylation and dephosphorylation of tau protein effects on microtubule interaction intracellular trafficking and neurodegeneration. *Biochem J*. 1997; 323(Pt 3): 577-591.
51. King ME, Gamblin TC, Kuret J, et al. Differential assembly of human tau isoforms in the presence of arachidonic acid. *J Neurochem*. 2000; 74(4): 1749-57.
52. Morales I, Jiménez JM, Mancilla M, et al. Tau oligomers and fibrils induce activation of microglial cells. *J Alzheimers Dis*. 2013; 37(4): 849-56.
53. Kochs G, Hummel R, Meyer D et al. Activation and substrate specificity of the human protein kinase C alpha and zeta isoenzymes. *Eur J Biochem*. 1993; 216: 597-606.
54. Mukai, H. The structure and function of PKN a protein kinase having a catalytic domain homologous to that of PKC. *J Biochem*. 2003; 133: 17-27.
55. Kawamata T, Taniguchi T, Mukai H, et al. A protein kinase PKN accumulates in Alzheimer neurofibrillary tangles and associated endoplasmic reticulum derived vesicles and phosphorylates tau protein. *J Neurosci*. 1998; 18(18): 7402-10.
56. Taniguchi T, Kawamata T, Mukai H, et al. Phosphorylation of tau is regulated by PKN. *J Biol Chem*. 2001; 276(13): 10025-31.
57. Rossner S, Mehlhorn G, Schliebs R, et al. Increased neuronal and glial expression of protein kinase C isoforms in neocortex of transgenic Tg2576 mice with amyloid pathology. *Eur J Neurosci*. 2001; 13(2): 269-78.
58. Miscia S, Ciccocioppo F, Lanuti P, et al. Abeta(1-42) stimulated T cells express P-PKC-delta and P-PKC-zeta in Alzheimer disease. *Neurobiol Aging*. 2009; 30(3): 394-406.
59. Zach S, Felk S, Gillardon F. Signal transduction protein array analysis links LRRK2 to Ste20 kinases and PKC zeta that modulate neuronal plasticity. *PLoS One*. 2010; 5: e13191.
60. Joshi YB, Giannopoulos PF, Chu J, et al. Absence of ALOX5 gene prevents stress-induced memory deficits synaptic dysfunction and tauopathy in a mouse model of Alzheimer's disease. *Hum Mol Genet*. 2014; 23(25): 6894-902.
61. Giannopoulos PF, Joshi YB, Chu J, et al. The 12-15-lipoxygenase is a modulator of Alzheimer's-related tau pathology in vivo. *Aging Cell*. 2013; 12(6): 1082-90.
62. Hosono T, Nishitsuji K, Nakamura T, et al. Arachidonic acid diet attenuates brain A β deposition in Tg2576 mice. *Brain Res*. 2015; 1613: 92-9.
63. Amtul Z, Uhrig M, Wang L, et al. Detrimental effects of arachidonic acid and its metabolites in cellular and mouse models of Alzheimer's disease, structural insight. *Neurobiol Aging*. 2012; 33(4): 831 e21-31.